arXiv:1512.01534v1 [math.RA] 4 Dec 2015

Oriented involutions and group identities on symmetric units of group algebras

Alexander Holguín-Villa *

2014-2015

Abstract

Let $\mathbb{F}G$ denote the group algebra of the locally finite group G over the infinite field $\mathbb{F}$ with $\text{char}(\mathbb{F}) \neq 2$, and let $\circledast : \mathbb{F}G \to \mathbb{F}G$ denote the involution defined by $\alpha = \Sigma\alpha_g g \mapsto \alpha^{\circledast} = \Sigma\alpha_g\sigma(g)g^*$, where $\sigma : G \to \{\pm 1\}$ is a group homomorphism (called an orientation) and $*$ is an involution of the group G. In this paper we prove, under some assumptions, that if the $\circledast$-symmetric units of $\mathbb{F}G$ satisfies a group identity then $\mathbb{F}G$ satisfies a polynomial identity, i.e., we give an affirmative answer to a Conjecture of B. Hartley in this setting. Moreover, in the case when the prime radical $\eta(\mathbb{F}G)$ of $\mathbb{F}G$ is nilpotent we characterize the groups for which the symmetric units $\mathcal{U}^+(\mathbb{F}G)$ do satisfy a group identity.

1 Introduction

Let $\mathbb{F}G$ denote the group algebra of the group G over of the field $\mathbb{F}$. Any involution $* : G \to G$ can be extended $\mathbb{F}$-linearly to an algebra involution of $\mathbb{F}G$. Such a map is called a group involution of $\mathbb{F}G$. A natural example is the so-called *classical involution*, which is induced from the map $g \mapsto g^* = g^{-1}$, for all $g \in G$.

Let $\sigma : G \to \{\pm 1\}$ be a non-trivial homomorphism (called an *orientation* of G). If $* : G \to G$ is a group involution, an *oriented group involution* of $\mathbb{F}G$ is defined by

$$\alpha = \sum_{g\in G} \alpha_g g \mapsto \alpha^{\circledast} = \sum_{g\in G} \alpha_g\sigma(g)g^*. \tag{1}$$

Notice that, as σ is non-trivial, $\text{char}(\mathbb{F})$ must be different from 2. It is clear that, $\alpha \mapsto \alpha^{\circledast}$ is an involution in $\mathbb{F}G$ if and only if $gg^* \in N = ker(\sigma)$ for all $g \in G$.

In the case when the involution on G is the classical involution, the map $\circledast$ is precisely the oriented involution introduced by S. P. Novikov, [16], in the context of K-theory.

We denote $\mathbb{F}G^+ = \{\alpha \in \mathbb{F}G : \alpha^{\circledast} = \alpha\}$ the set of symmetric elements of $\mathbb{F}G$ under $\circledast$ and, writing $\mathcal{U}(\mathbb{F}G)$ for the group of units of $\mathbb{F}G$, we let $\mathcal{U}^+(\mathbb{F}G)$ denote the set of symmetric units, i.e., $\mathcal{U}^+(\mathbb{F}G) = \{\alpha \in \mathcal{U}(\mathbb{F}G) : \alpha^{\circledast} = \alpha\}$.

A conjecture of Brian Hartley states that if the unit group $\mathcal{U}(\mathbb{F}G)$ of the group algebra $\mathbb{F}G$ of a torsion group G over a field $\mathbb{F}$ satisfies a group identity, then $\mathbb{F}G$ satisfies a polynomial identity. Let $\langle x_1, x_2, ...\rangle$ be the free group on a countable set of generators. A group H satisfies a group identity ($H \in$ GI or H is GI for short) if there exists a non-trivial reduced word $\omega(x_1, x_2, ..., x_n) \in \langle x_1, x_2, ...\rangle$ such that $\omega(h_1, h_2, ..., h_n) = 1$ for all $h_i \in H$.

*Partially supported by **CAPES** - Brazil.

We recall that a subset $S \subseteq \mathbb{F}G$ satisfies a polynomial identity ($S \in$ PI or S is PI for short) if there exits a non-zero polynomial $f(x_1, .., x_n)$ in the free associative algebra $\mathbb{F}\{x_1, x_2, ...\}$, in a countable set of non-commuting variables such that $f(a_1, ..., a_n) = 0$ for all $a_i \in S$. Group algebras satisfying a PI were classified by Passman and Isaacs-Passman, (see [17, Corollaries 3.8 and 3.10]).

Giambruno, Jespers and Valenti [4] solved the conjecture for semiprime group rings, and Giambruno, Sehgal and Valenti [8] solved it in general for group algebras over infinite fields. By using the results of [8], Passman [18] gave necessary and sufficient conditions for $\mathcal{U}(\mathbb{F}G)$ to satisfy a group identity, when $\mathbb{F}$ infinite. Subsequently, Liu [14] confirmed the conjecture for finite fields and Liu and Passman in [15] extended the results of [18] to this case. The same question for groups with elements of infinite order was studied by Giambruno, Sehgal and Valenti in [10]. For further details about these results, see G. Lee [13, Chapter 1].

Let $*$ be an involution of a group G extended linearly to $\mathbb{F}G$. Giambruno, Sehgal and Valenti [9] showed that if G is a torsion group, $\mathbb{F}$ is infinite with $\mathrm{char}(\mathbb{F}) \neq 2$, and $\mathcal{U}^+(\mathbb{F}G)$ satisfies a group identity then $\mathbb{F}G$ is PI. They also classified groups such that $\mathcal{U}^+(\mathbb{F}G)$ is *GI*, in the case of the classical involution.

Considering group involutions, Dooms and Ruiz [3] proved the following.

Theorem 1.1. ([3, Theorem 3.1]) *Let $\mathbb{F}$ be an infinite field with $\mathrm{char}(\mathbb{F}) \neq 2$ and let G be a non-abelian group such that $\mathbb{F}G$ is regular. Let $*$ be an involution on G. Suppose one of the following conditions holds:*

(i) *$\mathbb{F}$ is uncountable,*

(ii) *All finite non-abelian subgroups of G which are $*$-invariant have no simple components in their group algebra over $\mathbb{F}$ that are non-commutative division algebras other than quaternion algebras.*

Then $\mathcal{U}^+(\mathbb{F}G) \in GI$ if and only if G is an SLC-group with canonical involution. Moreover, in this case $\mathbb{F}G^+$ is a ring contained in $\zeta(\mathbb{F}G)$.

Using the last result and under some assumptions, Dooms and Ruiz proved that if $\mathcal{U}^+(\mathbb{F}G)$ satisfies a group identity then $\mathbb{F}G$ is PI, giving an affirmative answer to the Hartley's Conjecture in this setting. They also characterized the groups for which the symmetric units $\mathcal{U}^+(\mathbb{F}G)$ satisfy a group identity, when the prime radical $\eta(\mathbb{F}G)$ of $\mathbb{F}G$ is nilpotent. Giambruno, Polcino Milies and Sehgal [7] completely characterized group algebras of torsion groups, with group involutions such that $\mathcal{U}^+(\mathbb{F}G)$ is GI.

In this paper, we extend the results obtained by Dooms and Ruiz [3] to the case of oriented group involutions.

A group G is an LC-group (a group with "limited commutativity" property) if it is non-abelian and for any pair of elements $g, h \in G$, we have that $gh = hg$ if and only if at least one of g, h and gh is central in G. This family of groups was introduced by Goodaire. By Goodaire et al. [11, Proposition III.3.6], a group G is an LC-group with a unique non-trivial commutator s if and only if $G/\zeta(G) \cong C_2 \times C_2$, where C_2 is the cyclic group of order 2. If G is endowed with involution $*$, then we say that G is a special LC-group, or SLC-group, if it is an LC-group, it has a unique non-trivial commutator s and on such a group, the map $*$ is defined by

$$g^* = \begin{cases} g, & \text{if } g \text{ is central;} \\ sg, & \text{otherwise.} \end{cases} \tag{2}$$

We refer to this as the *canonical* involution on an SLC-group. For instance, if $*$ is the classical involution, then it is easy to see that the SLC-groups are precisely the Hamiltonian 2-groups.

Throughout this paper $\mathbb{F}$ will always denote an infinite field with $\mathrm{char}(\mathbb{F}) \neq 2$, G a group, $*$ and σ an involution and a non-trivial orientation of G, respectively. $\circledast$ will denote an oriented group involution of $\mathbb{F}G$ given by expression (1).

2 Preliminaries and notations

Let R be a ring with involution $\star$. Let $\mathcal{U}(R)$ its group of units and $\mathcal{U}^+(R) = \mathcal{U}(R) \cap R^+$ the set of symmetric units. It is well-known that central idempotents are very important in the study of group identities. Moreover the following fact is proved:

Lemma 2.1. ([9, Theorem 2]) *Let R be a semiprime ring with involution $\star$ such that $\mathcal{U}^+(R)$ is GI. Then every symmetric idempotent of R is central.*

For a given prime p, an element $x \in G$ will be a called a p-element if its order is a power of p and it is called p'-element if its order is finite and, not divisible by p. Moreover, a torsion subgroup H of G is a p'-subgroup if every element $h \in H$ is a p'-element. We agree that if $p = 0$ every torsion subgroup is a p'-subgroup.

An immediate consequence in the setting of group algebras of the Lemma 2.1 is the following: Let $\mathbb{F}$ be a field with $\mathrm{char}(\mathbb{F}) \geq 0$ and G a group such that $\mathbb{F}G$ is semiprime. If $\mathcal{U}^+(\mathbb{F}G)$ is GI under classical involution, then every torsion p'-subgroup of G is normal in G. This fact give important information on cyclic subgroups, a property which is lost for a group involutions $*$, extended linearly to the group algebra $\mathbb{F}G$. Semisimple algebras whose units satisfy a GI were widely studied, see for instance [5, Theorem 1], [3, Theorem 2.2], [1, Theorem 3.1] and [7, Lemma 2.1].

The following three lemmas will be needed in the sequel:

Lemma 2.2. ([7, Lemma 2.1]) *Let R be a finite dimensional semisimple algebra with involution $\star$ over infinite field K,* $\mathrm{char}(K) \neq 2$. *Suppose that $\mathcal{U}^+(R)$ is GI. Then R is a direct sum of simple algebras of dimension at most four over their centers and the symmetric elements R^+ are central in R, i.e.,*

$$R \cong D_1 \oplus D_2 \oplus ... \oplus D_k \oplus M_2(\mathbb{F}_1) \oplus M_2(\mathbb{F}_2) \oplus ... \oplus M_2(\mathbb{F}_l).$$

Lemma 2.3. ([3, Theorem 2.2]) *Let R be a semisimple K-algebra with involution $\star$, where K is an infinite field with* $\mathrm{char}(K) \neq 2$. *Suppose one of the following conditions holds:*

(i) *K is uncountable,*

(ii) *R has no simple components that are non-commutative division algebras other than quaternion algebras.*

Then $\mathcal{U}^+(R) \in GI$ if and only if R^+ is central in R.

Lemma 2.4. ([13, Lemma 2.3.5]) *Suppose that R is an K-algebra with involution $\star$, where* $\mathrm{char}(K) \neq 2$. *Let I be a nil ideal $\star$-invariant. If $\mathcal{U}^+(R) \in GI$, then $\mathcal{U}^+(R/I) \in GI$. Conversely, if $p > 0$, I is nil of bounded exponent and $\mathcal{U}^+(R/I) \in GI$, then $\mathcal{U}^+(R) \in GI$.*

Remark 1. *By [6, Lemma 2.4], under the assumptions of Lemma 2.3, it follows that $\mathcal{U}^+(R) \in GI$ is equivalent to R^+ is Lie n-Engel, for some n.*

We conclude this section with a result due to Jespers an Ruiz Marín [12], where the SLC groups arise naturally.

Lemma 2.5. ([12, Theorem 2.4]) *Let $\mathbb{F}$ be a field with $\mathrm{char}(\mathbb{F}) \neq 2$ and let G be a group with an involution $*$ extended $\mathbb{F}$-linearly to $\mathbb{F}G$. Then $\mathbb{F}G^+$ is commutative if and only if G is abelian or an SLC group. In this case, $\mathbb{F}G^+ = \zeta(\mathbb{F}G)$.*

3 Results

We need the following results about group rings with an oriented group involution:

Lemma 3.1. ([2, Lemma 1.1]) *Let R be a commutative ring with unity of characteristic different from 2 and let G be a group with a non-trivial orientation σ and an involution $*$. Suppose that RG^+ is commutative under oriented group involution and let $g \in (G \backslash N) \backslash G^+$, $h \in G$. Then one of the following holds:*

(i) *$gh = hg$; or*

(ii) *$\mathrm{char}(R) = 4$ and $gh = g^*h^* = hg^* = h^*g$.*

Lemma 3.2. ([2, Theorem 2.2]) *Let R be a commutative ring with unity of characteristic different from 2 and let G be a non-abelian group with a non-trivial orientation σ and an involution $*$. Then, RG^+ is a commutative ring if and only if one of the following conditions holds:*

(i) *$N = ker(\sigma)$ is an abelian group and $(G \setminus N) \subset G^+$;*

(ii) *G and N have the* LC*-property, and there exists a unique non-trivial commutator s such that the involution $*$ is given by*

$$g^* = \begin{cases} g, & \text{if } g \in N \cap \zeta(G) \text{ or } g \in (G \setminus N) \setminus \zeta(G); \\ sg, & \text{if otherwise.} \end{cases}$$

(iii) *$\mathrm{char}(R) = 4$, G has the* LC*-property, and there exists a unique non-trivial commutator s such that the involution $*$ is the canonical involution.*

We will denote by $(\alpha, \beta) = \alpha^{-1}\beta^{-1}\alpha\beta$ the multiplicative commutator of $\alpha, \beta \in \mathcal{U}(\mathbb{F}G)$. Recursively, one defines $(\alpha_1, \alpha_2, \ldots, \alpha_n) = ((\alpha_1, \alpha_2, \ldots, \alpha_{n-1}), \alpha_n)$ where $\alpha_1, \alpha_2, \ldots, \alpha_n \in \mathcal{U}(\mathbb{F}G)$.

Recall that a ring R with identity is said to be (von Neumann) regular if for any $x \in R$ there exists an $y \in R$ such that $xyx = x$. Villamayor [19] showed that the group algebra $\mathbb{F}G$ is regular if and only if G is locally finite and has no elements of order p in case $\mathbb{F}$ has characteristic $p > 0$. Note that in this case $\mathbb{F}G$ is semiprime, [17, Theorem 4.2.13] and the set of p-elements P is trivial (in case $\mathrm{char}(\mathbb{F}) = 0$, we agree that that $P = 1$).

We are now able to classify the groups with a regular group algebra over an infinite field $\mathbb{F}$ with $\mathrm{char}(\mathbb{F}) \neq 2$ for which the $\circledast$-symmetric units satisfy a GI.

Theorem 3.1. *Let $\mathbb{F}$ be an infinite field with $\mathrm{char}(\mathbb{F}) \neq 2$ and let G be a non-abelian group such that $\mathbb{F}G$ is regular. Let $\sigma : G \to \{\pm 1\}$ be a non-trivial orientation and an involution $*$ on G. Suppose one of the following conditions holds:*

(i) $\mathbb{F}$ *is uncountable,*

(ii) *All finite non-abelian subgroups of* G *which are* $*$*-invariant have no simple components in their group algebra over* $\mathbb{F}$ *that are non-commutative division algebras other than quaternion algebras.*

Then $\mathcal{U}^+(\mathbb{F}G) \in GI$ *if and only if one of the following conditions holds:*

1. $N = ker(\sigma)$ *is an abelian group and* $(G \setminus N) \subset G^+$*;*
2. G *and* N *have the* LC*-property, and there exists a unique non-trivial commutator* s *such that the involution* $*$ *is given by*

$$g^* = \begin{cases} g, & \text{if } g \in N \cap \zeta(G) \text{ or } g \in (G \setminus N) \setminus \zeta(G); \\ sg, & \text{if otherwise.} \end{cases} \tag{3}$$

Consequently, $\mathcal{U}^+(\mathbb{F}G) \in GI$ *if and only if* $\mathcal{U}^+(\mathbb{F}G)$ *is an abelian group.*

Proof. Assume that $\mathcal{U}^+(\mathbb{F}G) \in$ GI and let $N = ker(\sigma)$. Then $\mathcal{U}^+(\mathbb{F}N) \in$ GI. Hence, by Theorem 1.1, we have two possibilities for N; either

(A) N is an abelian group; or

(B) N has the LC-property, and there exists a unique non-trivial commutator s such that the involution $*$ in N, is the canonical involution.

Now, let $g, h \in G \setminus \zeta(G)$. Consider the collection $\{H_i\}_{i \geq 1}$ of all finite subgroups of G which are $*$-invariant and contain g and h. It is clear that $G = \bigcup_i H_i$ and that $\mathcal{U}^+(\mathbb{F}H_i) \in$ GI. Since $\mathbb{F}G$ is regular, we have that all $\mathbb{F}H_i$ are semisimple.

Set $\sigma_i = \sigma|_{H_i}$ and let $N_i = ker(\sigma_i)$. By Lemma 2.3, $\mathbb{F}H_i^+$ is central in $\mathbb{F}H_i$ for all i, and applying the Lemma 3.2, one of the following conditions holds:

(a) N_i is an abelian group and $(H_i \setminus N_i) \subset H_i^+$; or

(b) H_i and N_i have the LC-property, and there exists a unique non-trivial commutator s such that the involution $*$ is as given in the statement.

If s is not a unique non-trivial commutator of G, then there exist $x, y \in G$ such that $(x, y) \neq s$. But $x, y \in H_i$, for some i, for instance, $H_i = \langle x, y, g, h, x^*, y^*, g^*, h^* \rangle$. Therefore $(x, y) = (g, h) = s$, a contradiction.

It is easy to see that, $N = ker(\sigma) = \bigcup_i N_i = \bigcup_i ker(\sigma_i)$ and $\bigcup_i (H_i \setminus N_i) = (G \setminus N)$ and hence we have (1).

Claim: For all i such that $(H_i \setminus N_i) \not\subseteq H_i^+$, $\zeta(N_i) = N_i \cap \zeta(H_i)$. In fact, we have showed above that for all i, $\mathbb{F}H_i^+$ is a commutative ring. Let $h \in H_i \setminus N_i$. If h is not central then, there exists $g \in H_i$ such that $hg \neq gh$ and by Lemma 3.1, it follows that $h \in H_i^+$. If h is central, take $x \in N_i \setminus \zeta(N_i)$. Then $xh \in (H_i \setminus N_i) \setminus \zeta(H_i)$ and again, as above $xh \in H_i^+$. Therefore, $xh = (xh)^* = h^*x^* = sh^*x$, i.e., $h^* = sh$.

Now, let $x \in \zeta(N_i) \setminus \zeta(H_i)$. Then, there exists $y \in H_i \setminus N_i$ such that $xy \neq yx$ and $y^* = y$. Since $\mathbb{F}N_i^+$ is commutative, for all i, we have that $x^* = x$. Thus $xy \in (H_i \setminus N_i) \setminus \zeta(H_i)$ and again, by Lemma 3.1, $xy \in H_i^+$. Therefore $xy = (xy)^* = y^*x^* = yx$, a contradiction. Hence $\zeta(N_i) = N_i \cap \zeta(H_i)$.

Condition (b) above implies that for all i, $H_i^+ = \zeta(N_i) \cup [(H_i \setminus N_i) \setminus \zeta(H_i)]$. Now, it is clear that $\zeta(G) \subseteq \bigcup_i \zeta(H_i)$. When $x \in \bigcup_i \zeta(H_i)$, then $x \in \zeta(H_j)$ for some j and hence, $xg = gx$ where $g \in G \setminus \zeta(G)$ is arbitrary, i.e., $x \in \zeta(G)$ and $\zeta(G) = \bigcup_i \zeta(H_i)$.
Therefore $* : G \to G$ is as given in the statement, and (2) holds.

The converse is clear, because conditions (1) and (2) by Lemma 3.2 imply that $\mathbb{F}G^+$ is commutative and hence, $(u, v) = 1$ is a GI for $\mathcal{U}^+(\mathbb{F}G)$.

The last assertion is now clear.

□

The next lemma which we need is proved in [6, Lemma 2.6].

Lemma 3.3. *Let $\mathbb{F}$ be a field with $\mathrm{char}(\mathbb{F}) = p > 2$, G a finite group and $\mathcal{J}$ the Jacobson radical of $\mathbb{F}G$. Suppose that $\mathbb{F}G/\mathcal{J}$ is isomorphic to a direct sum of simple algebras of dimension at most four over their centers. Then the set P of p-elements of G is a subgroup.*

To handle group algebras which are not necessarily regular, we need the following two lemmas which are the natural extensions of known results. As usual, for a normal subgroup H of G we denote by $\Delta(G, H)$ the kernel of the map $\mathbb{F}G \xrightarrow{\Psi} \mathbb{F}(G/H)$ defined by

$$\sum_{g \in G} \alpha_g g \mapsto \sum_{g \in G} \alpha_g gH$$

and $\Delta(G, G) = \Delta(G)$ is the augmentation ideal.

Lemma 3.4. *Let G be a locally finite group and $\mathrm{char}(\mathbb{F}) = p \neq 2$. If $\mathcal{U}^+(\mathbb{F}G) \in GI$, then the set P of p-elements of G is a subgroup.*

Proof. Let $g, h \in P$ and let $H = \langle g, h, g^*, h^* \rangle$. Since G is locally finite, then H is finite. Moreover, H is $*$-invariant and $H \subset N = ker(\sigma)$ (every element $x \in H$ has odd order). Since $\mathbb{F}H$ is a finite dimensional algebra the Jacobson radical $\mathcal{J}$ is nilpotent. Let $R = \mathbb{F}H/\mathcal{J}$. Then R is semisimple and, by Lemma 2.4, $\mathcal{U}^+(R)$ satisfies a group identity. Hence by Lemma 2.2, R is a direct sum of simple algebras of dimension at most four over their centers. Finally, the lemma above shows that P is a subgroup.

□

Lemma 3.5. *Let $\mathbb{F}$ be a field with $\mathrm{char}(\mathbb{F}) = p > 2$ and G a group such that $\mathcal{U}^+(\mathbb{F}G)$ satisfies a GI $\omega(x_1, ..., x_n) = 1$, under an oriented group involution $\circledast$. If H is a normal $*$-invariant p-subgroup of G, and either H is finite or G is locally finite, then $\mathcal{U}^+(\mathbb{F}(G/H))$ satisfies $\omega(x_1, ..., x_n) = 1$.*

Proof. Since H is a p-subgroup, then $H \subset N$ and hence H is $\circledast$-invariant.
If H is finite, then by [13, Lemma 1.1.1], $\Delta(G, H)$ is nilpotent. If G is locally finite, then for every n-tuple $\alpha_1, ..., \alpha_n$ in $\mathbb{F}G$, set $G_1 = \langle Supp(\alpha_i), Supp((\alpha_i)^{-1}) \rangle$ and taking $H_1 = G_1 \cap H$ we have again by [13, Lemma 1.1.1] that $\Delta(G, H_1)$ is nilpotent.

Now $\mathbb{F}(G/H) \cong \mathbb{F}G/\Delta(G, H)$ and the epimorphism $\mathbb{F}G \to \mathbb{F}(G/H)$ induces an epimorphism $\mathcal{U}^+(\mathbb{F}G) \to \mathcal{U}^+(\mathbb{F}(G/H))$. By Lemma 2.4, we have that $\mathcal{U}^+(\mathbb{F}(G/H))$ satisfies a GI.

□

Now we obtain a result about group algebras which are not regular:

Theorem 3.2. *Let $g \mapsto g^*$ be an involution on a locally finite group G, $\sigma : G \to \{\pm 1\}$ a non-trivial orientation with $N = ker(\sigma)$ and $\mathbb{F}$ an infinite field with $\mathrm{char}(\mathbb{F}) = p \neq 2$. Suppose that $\mathcal{U}^+(\mathbb{F}G) \in GI$ and that one of the following conditions holds:*

(i) $\mathbb{F}$ *is uncountable,*

(ii) *All finite non-abelian subgroups of* G/P *which are* $*$*-invariant have no simple components in their group algebra over* $\mathbb{F}$ *that are non-commutative division algebras other than quaternion algebras,*

then we have that

1. $\overline{G} = G/P$ *is abelian, or*
2. $\overline{N} = N/P = ker(\overline{\sigma})$ *is abelian and* $(\overline{G} \setminus \overline{N}) \subset \overline{G^+}$*, or*
3. $\overline{G}$ *and* $\overline{N}$ *have the* LC*-property and there exists a unique non-trivial commutator* $\overline{s}$ *such that the involution* $\overline{*}$ *in* $\overline{G}$ *is given by*

$$\overline{g^*} = \begin{cases} \overline{g}, & \text{if } \overline{g} \in \overline{N} \cap \zeta(\overline{G}) \text{ or } \overline{g} \in (\overline{G} \setminus \overline{N}) \setminus \zeta(\overline{G}); \\ \overline{sg}, & \text{if otherwise.} \end{cases} \tag{4}$$

Moreover, $\mathbb{F}G \in PI$.

Proof. By Lemma 3.4, we have that P is a normal subgroup of G and by Lemma 3.5 $\mathcal{U}^+(\mathbb{F}(G/P))$ is GI. Since $\mathbb{F}(G/P)$ is regular, it follows that either G/P is abelian, or $\overline{N}$ and $\overline{G}$ satisfy one of the conditions of Theorem 3.1 and the involution $\overline{*} : \overline{G} \to \overline{G}$ is as given in the statement. Moreover, $\mathbb{F}(G/P)^+$ is central in $\mathbb{F}(G/P)$ and thus $\mathbb{F}(G/P)$ is PI. Since $\mathbb{F}(G/P) \cong \mathbb{F}G/\Delta(G,P)$ and $\Delta(G,P)$ is a nil subring of $\mathbb{F}G$ $\circledast$-invariant, by [9, Remark 2] we have that $\Delta(G,P)$ is PI. Hence, $\mathbb{F}G$ is also PI.

□

To obtain sufficient conditions for locally finite groups G such that $\mathcal{U}^+(\mathbb{F}G) \in$ GI, we need the following lemma:

Lemma 3.6. *Let* $\mathbb{F}$ *be a field with* $\mathrm{char}(\mathbb{F}) = p \neq 2$*. Let* G *be a locally finite group with an involution* $*$ *and a non-trivial orientation* σ*. If* P *is a subgroup of bounded exponent, and either* G/P *is abelian, or* G/P *and* N/P *are as in Theorem 3.1, then* $\mathcal{U}^+(\mathbb{F}G) \in GI$*.*

Proof. Suppose P is a subgroup of bounded exponent and that N/P, G/P, $*$ and σ are as in the statement. Then by Lemma 3.2, $\mathcal{U}^+(\mathbb{F}\overline{G})$ is abelian. Hence $(\mathcal{U}^+(\mathbb{F}G), \mathcal{U}^+(\mathbb{F}G)) \subset 1 + \Delta(G,P)$. Now $\Delta(G,P)$ is nil of bounded exponent and thus $(\mathcal{U}^+(\mathbb{F}G), \mathcal{U}^+(\mathbb{F}G))^{p^n} = 1$ for some $n \geq 0$. Hence $\mathcal{U}^+(\mathbb{F}G) \in$ GI.

□

For $g \in G$, let $C_G(g) = \{h \in G : hg = gh\}$ be the centralizer of g in G. Set $\Phi(G) = \{g \in G : [G : C_G(g)] < \infty\}$ the finite conjugacy subgroup of G and $\Phi_p(G) = \langle P \cap \Phi(G) \rangle$.

Theorem 3.3. *Let* $g \mapsto g^*$ *be an involution on a locally finite group* G*,* $\sigma : G \to \{\pm 1\}$ *a non-trivial orientation and* $\mathbb{F}$ *an infinite field with* $\mathrm{char}(\mathbb{F}) = p \neq 2$*. Suppose that the prime radical* $\eta(\mathbb{F}G)$ *of* $\mathbb{F}G$ *is a nilpotent ideal and that one of the following conditions holds:*

(i) $\mathbb{F}$ *is uncountable,*

(ii) *All finite non-abelian subgroups of* G/P *which are* $*$*-invariant have no simple components in their group algebra over* $\mathbb{F}$ *that are non-commutative division algebras other than quaternion algebras.*

Then $\mathcal{U}^+(\mathbb{F}G) \in GI$ if and only if P is a finite normal subgroup and G/P is abelian or G/P and N/P are as in the Theorem 3.1.

Proof. Suppose that $\mathcal{U}^+(\mathbb{F}G) \in$ GI, then by Lemma 3.4 we have that P is a normal subgroup. Now, by Theorem 3.2, either G/P is abelian or G/P and N/P are as in Theorem 3.1 and hence, $\mathbb{F}G \in$ PI. Thus by [18, Theorem 5.2.14], $[G : \Phi(G)] < \infty$ e $|\Phi'(G)| < \infty$. Since $\eta(\mathbb{F}G)$ is nilpotent [18, Theorem 8.1.12] gives that $\Phi_p(G) = P \cap \Phi(G)$ is a finite normal p-subgroup. As $P\Phi(G)/\Phi(G) \cong P/P \cap \Phi(G)$ is finite, then P is finite.

Now, the converse is clear by Lemma 3.6.

□

As a corollary following the arguments in Dooms and Ruiz [3, Corollary 3.7], we obtain a characterization of the locally finite groups with semiprime group algebra such that the set of $\circledast$-symmetric units is GI.

Corollary 3.1. *Let $g \mapsto g^*$ be an involution on a locally finite group G, $\sigma : G \to \{\pm 1\}$ a non-trivial orientation and $\mathbb{F}$ an infinite field with* $\mathrm{char}(\mathbb{F}) = p \neq 2$ *such that $\mathbb{F}G$ is semiprime. Suppose one of the following conditions holds:*

(i) *$\mathbb{F}$ is uncountable,*

(ii) *All finite non-abelian subgroups of G/P which are $*$-invariant have no simple components in their group algebra over $\mathbb{F}$ that are non-commutative division algebras other than quaternion algebras.*

Then $\mathcal{U}^+(\mathbb{F}G) \in GI$ if and only if either G/P is abelian or G/P and N/P are as in Theorem 3.1.

Proof. Suppose $\mathcal{U}^+(\mathbb{F}G) \in$ GI, then following the lines of the proof of Theorem 3.3, $\mathbb{F}G$ is semiprime PI. Hence by [18, Theorem 4.2.13] $\Phi_p(G) = 1$ and by previous proof, we get that $P = 1$. Thus $\mathbb{F}G$ is regular, and the result follows from Theorem 3.1.

□

References

[1] O. Broche Cristo, A. Dooms and M. Ruiz. *Unitary units satisfying a group identity*, Comm. Algebra **37** (2009):1729-1738.

[2] O. Broche Cristo, C. Polcino Milies, *Symmetric elements under oriented involutions in group rings*, Comm. Algebra **34** (2006):3347-3356.

[3] A. Dooms and M. Ruiz. *Symmetric units satisfying a group identity*, J. Algebra **308** (2007):742-750.

[4] A. Giambruno, E. Jespers and A. Valenti. *Group identities on units of rings*, Arch. Math. **63** (1994):291-296.

[5] A. Giambruno and C. Polcino Milies. *Unitary units and skew elements in group algebras*, Manuscripta Math. **111** (2003):195-209.

[6] A. Giambruno, C. Polcino Milies, S. K. Sehgal, *Lie properties of symmetric elements in group rings*, J. Algebra **321** (2009):890-902.

[7] A. Giambruno, C. Polcino Milies and S. K. Sehgal. *Group identities on symmetric units*, J. Algebra **322** (2009):2801-2815.

[8] A. Giambruno, S. K. Sehgal and A. Valenti. *Group algebras whose units satisfy a group identity*, Proc. Amer. Math. Soc. **125** (1997):629-634.

[9] A. Giambruno, S. K. Sehgal and A. Valenti. *Symmetric units and group identities*, Manuscripta Math. **96** (1998):443-461.

[10] A. Giambruno, S. K. Sehgal and A. Valenti. *Group identities on units of group algebras*, J. Algebra **226** (2000):488-504.

[11] E. G. Goodaire, E. Jespers and C. Polcino Milies. *Alternative loop rings*, North-Holland, Amsterdam (1996).

[12] E. Jespers, M. Ruiz Marín. *On symmetric elements and symmetric units in group rings*, Comm. Algebra **34** (2006):727-736.

[13] G. T. Lee. *Group identities on units and symmetric units of group rings*, Algebra and Applications, vol. 12, Springer-Verlag, London, (2010).

[14] C. H. Liu. *Group algebras whit units satisfying a group identity*, Proc. Amer. Math. Soc. **127** (1999):327-336.

[15] C. H. Liu, D. S. Passman. *Group algebras whit units satisfying a group identity II*, Proc. Amer. Math. Soc. **127** (1999):337-341.

[16] S. P. Novikov. *Algebraic construction and properties of Hermitian analogs of K-theory over rings with involution from the viewpoint of Hamiltonian formalism. Applications to differential topology and the theory of characteristic classes. I. II.* Izv. Akad. Nauk SSSR Ser. Mat. **34** (1970):253-288; ibid **34** (1970):475-500.

[17] D. S. Passman. *The algebraic structure of group rings*, Pure and Applied Mathematics, Wiley-Interscience [John Wiley & Sons], New York, (1977).

[18] D. S. Passman. *Group algebras whose units satisfy a group identity II*, Proc. Amer. Math. Soc. **125** (1997):657-662.

[19] E. Villamayor. *On weak dimension of algebras*, Pac. J. Math. **9** (15959):941-951.

Alexander Holguín Villa
Instituto de Matemática e Estatística
Universidade de São Paulo, SP, Brazil
Rua do Matão, 1010 - Cidade Universitária
CP 66281- CEP 05508-090
E-mail: alexholguinvilla@gmail.com